

## N O T I C E

THIS DOCUMENT HAS BEEN REPRODUCED FROM  
MICROFICHE. ALTHOUGH IT IS RECOGNIZED THAT  
CERTAIN PORTIONS ARE ILLEGIBLE, IT IS BEING RELEASED  
IN THE INTEREST OF MAKING AVAILABLE AS MUCH  
INFORMATION AS POSSIBLE

(NASA-TM-80679) INITIAL GEOMAGNETIC FIELD  
MODEL FROM MAGSAT (NASA) 19 p HC A02/MF A01  
CSCL 04A

N80-25967

Unclas  
G3/46 22942



## Technical Memorandum 80679


# Initial Geomagnetic Field Model from MAGSAT

R. A. Langel, R. H. Estes, G. D. Mead,  
E. B. Fabiano and E. R. Lancaster

APRIL 1980

National Aeronautics and  
Space Administration

Goddard Space Flight Center  
Greenbelt, Maryland 20771



INITIAL GEOMAGNETIC FIELD MODEL FROM MAGSAT

R.A. Langel<sup>1</sup>  
R.H. Estes<sup>2</sup>  
G.D. Mead<sup>1</sup>  
E.B. Fabiano<sup>3</sup>  
E.R. Lancaster<sup>1</sup>

April 1980

Submitted to Geophysical Research Letters

<sup>1</sup>Goddard Space Flight Center, Greenbelt, MD 20771

<sup>2</sup>Business and Technological Systems, Seabrook, MD 20801

<sup>3</sup>U.S. Geological Survey, Denver, CO 80225

### ABSTRACT

Magsat data from the magnetically quiet days of November 5-6, 1979, were used to derive a thirteenth degree and order spherical harmonic geomagnetic field model, MGST(3/80). The model utilized both scalar and vector data and fit that data with standard deviations of 8, 52, 55 and 97 nT for the scalar magnitude,  $B_r$ ,  $B_\theta$ , and  $B_\phi$ , respectively. When compared with earlier models, the earth's dipole moment continues to decrease at a rate of about 26 nT/yr. Evaluation of earlier models with Magsat data shows that the scalar field at the Magsat epoch is best predicted by the POGO(2/72) model but that the AWC/75 and IGS/75 are better for predicting vector fields.

## INTRODUCTION

The NASA spacecraft Magsat (NASA/GSFC, 1979) was successfully launched into a low-altitude, near-polar orbit on October 30, 1979. Intended specifically to map the magnetic field near the Earth (Langel et al., 1977; Langel, 1979), Magsat's instrumentation includes a Cesium vapor scalar magnetometer of accuracy  $\pm 1$  nT (nanotesla) and a fluxgate magnetometer of accuracy better than 3 nT rms (Acuna et al., 1978). The scalar magnetometer has been successfully utilized to calibrate the vector magnetometer to an rms difference in field magnitude of less than 1 nT for the data reported herein. Initial attitude determination utilized an Earth horizon scanner and a precision sun sensor. The resulting accuracy is estimated to be somewhat better than 20 arc-minutes rms, which is equivalent to a component accuracy of 290 nT in a 50000 nT field or 175 nT in a 30000 nT field. Later data processing (Langel and Berbert, 1979) will utilize on-board star cameras and an optical system to measure the attitude of the vector magnetometer relative to the star cameras. This is expected to improve the attitude accuracy to about 20 arc-seconds rms, with corresponding component accuracies of 3-6 nT. The accuracy of orbit determination is better than the 60 m radial and 300 m horizontal accuracy required for a 3 nT rms overall accuracy of the scalar data (Langel, 1976). For the initial attitude accuracy, orbital error has negligible effect on the vector data.

The anticipated lifetime at launch was five months. However, it now appears that orbital decay is considerably less than predicted and we expect a seven to seven-and-a-half month lifetime before reentry.

Initial orbital parameters for Magsat are:

Apogee:	561.2 km
Perigee:	352.4 km
Inclination:	$96.76^{\circ}$ .

Several magnetically quiet days of Magsat data have now become available and are suitable for deriving a preliminary field model and for evaluating existing geomagnetic field models for the epoch of November 1979.

#### THE MAGSAT MODEL

Of the data currently available to us for analysis, November 5 and 6 have the lowest Kp indices. The highest three-hourly Kp value is 1+, which occurred only once. Plots of Dst and of the data itself also indicated that these days were indeed very quiet. Accordingly, a selection of 7468 data points from these days was used to derive an initial Magsat model whose spherical harmonic coefficients are given in Table 1. This included 2414 scalar points whose distribution is depicted in Figure 1. Global coverage was excellent; all but 10 of  $648 \times 10^{\circ} \times 10^{\circ}$  blocks contained at least one data point. Also included were 1679 values of  $B_r$ , 1681 values of  $B_{\theta}$ , and 1694 values of  $B_{\phi}$ , where  $B_r$ ,  $B_{\theta}$  and  $B_{\phi}$  are the field components measured in spherical coordinates, for latitudes below  $50^{\circ}$ . Their distribution for those latitudes is similar to that of the scalar data shown in Figure 1. Vector data above  $50^{\circ}$  latitude were not included because of non-curl-free fields from field-aligned currents. Such fields are transverse to the main field and, as shown by Langel (1974), have little or no effect on the amplitude of the scalar measurements. All data are at twilight local times because of the orientation of the Magsat orbit.

The fitting method was essentially that described by Cain et al. (1967).

A degree and order thirteen model was chosen because it was justified by the density of the data and because studies by Phillips (unpublished, personal communication), and Cain (1976) indicate that a thirteenth degree and order representation is sufficient to represent the main field of the Earth. Any such choice is, of course, somewhat subjective.

This model fits the selected data with the mean and standard deviations shown in the following table.

<u>COMPONENTS</u>	<u>MEAN DEVIATION (nT)</u>	<u>STANDARD DEVIATION (nT)</u>
Scalar Magnitude	0	8
$B_r$	22	52
$B_\theta$	10	55
$B_\phi$	84	97

The source of the non-zero means for the vector data appears to be a bias of 4-7 arc-minutes in the initial attitude solution. The standard deviation of the scalar data is mainly due to a combination of fields from external and crustal sources, neither of which is modeled. The standard deviation of the vector data is due to current lack of precision in spacecraft attitude.

Evaluation of such a model is best done by comparison with independent data. Although a global set of independent data is not now available to us, comparisons have been made with a set of 44 repeat station measurements obtained by the U.S. Geological Survey in 1978 and 1979. The root mean square (rms) of residuals of total intensity at these stations was determined to be of the order of 170 nT, which may be considered as a good fit to the surface data. We have also plotted residuals of several additional days of Magsat

data relative to this model. These plots indicate that the model is a good representation of the quiet-time field at the spacecraft.

The model was used to compute the field magnitude B, declination D, horizontal field H and vertical field Z at 400 km altitude. Contour plots are shown in Figure 2.

No secular variation terms are included in the model. However, the usefulness of the model could be extended in time by including secular variation terms from models such as IGS/75, AWC/75, or IGRF 1975 (whose terms consist of averages of IGS/75 and AWC/75).

#### STRENGTH OF THE CENTERED DIPOLE

The first approximation of the geomagnetic field is that of the field of a dipole located at the Earth's center and inclined to its axis of rotation. The strength of that dipole is measured by  $MR^{-3}$ , where  $M$  is the dipole moment computed from the first three harmonic coefficients and  $R$  is the mean radius of the Earth.  $MR^{-3}$  has been decreasing for some years (Harwood and Malin, 1976; Barraclough et al., 1978), resulting in speculation that we may be observing a reversal of the Earth's field. We have computed  $MR^{-3}$  from the new model and compared it to earlier models, as summarized in Table 2 and Figure 3. The Magsat results show a continuing decrease of  $MR^{-3}$ , consistent with earlier results. A comparison of the dipole moment measured by Magsat with that measured by POGO around 1968 indicates a rate of decrease of 26 nT/year. The slope of the line in Figure 3 is -27 nT/year. These rates are somewhat larger than the 10-20 nT/year rates measured in the 1940-1960 time frame (Harwood and Malin, 1976). If the higher rates continue, the Earth's field would reverse in about 1200 years.



### EVALUATION OF EARLIER MODELS

The availability of Magsat data makes it convenient to evaluate the predictive capability of those field models currently in common use in the scientific community (see also Mead, 1979). For this evaluation 4825 scalar points and 6637 vector points at all latitudes were selected from November 5 and 6, 1979. The best known and perhaps most widely used of the current models is the IGRF (International Geomagnetic Reference Field, Zmuda, 1971; updated for use from 1975, IAGA, 1976). Other models chosen for evaluation were IGS/75 (Barracough et al., 1975), AWC/75 (Peddie and Fabiano, 1976) and POGO(2/72) (Langel et al., 1980). Table 3 shows the mean and standard deviation of the selected Magsat data with respect to each of these models.

The POGO(2/72) model was derived from satellite scalar field data only. Its predictive capability for the scalar field at the Magsat epoch is, as might be expected, better than the other models shown in Table 3. For component prediction the AWC/75 and IGS/75 models are very close in capability, with perhaps a slight edge to IGS/75. The IGRF 1975 model is significantly poorer in its predictions of the vector field at the Magsat epoch.

### SUMMARY

A global set of scalar and vector data from Magsat was used to construct a thirteenth degree and order geomagnetic field model at epoch 1979.85. The model fits the Magsat scalar data with a standard deviation of 8 nT, and also provides a good fit to 44 recent repeat station measurements. Comparison with previous models indicates that the Earth's dipole moment continues to decrease, at a rate of 20 nT/year. Evaluation of earlier models with Magsat data shows that the POGO(2/72) model gave the best predictions of the scalar field, but that the IGS/75 and AWC/75 models were better at predicting the vector components.

ACKNOWLEDGEMENTS

Special thanks go to Gil Ousley and his staff at GSFC and to L.D. Eckard and his staff at Johns Hopkins University Applied Physics Laboratory for successfully building and flying the Magsat satellite, to M. Acuna and W.H. Farthing of GSFC for providing the Magsat magnetometers, to John Berbert, Earl Beard and associates at GSFC and Don Berman and staff at Computer Sciences Corporation for their data processing efforts and to Jim Strayer of Business and Technological Systems for long hours at the computer.

## REFERENCES

- Acuna, M.H., C.S. Searce, J.B. Seek and J. Scheifele, The Magsat Vector Magnetometer - A Precision Fluxgate Magnetometer for the Measurement of the Geomagnetic Field, NASA TM-79656, October 1978.
- Barracough, D.R., J.M. Harwood, B.R. Leaton, and S.R.C. Malin, A Model of the Geomagnetic Field at Epoch 1975, Geophys. J. Royal Astr. Soc., 43, 645-659, 1975.
- Barracough, D.R., J.M. Harwood, B.R. Leaton, and S.R.C. Malin, A Definitive Model of the Geomagnetic Field and its Secular Variation for 1965 - I. Derivation of Model and Comparison with the IGRF, Geophys. J. Royal Astr. Soc., 55, 111-121, 1978.
- Cain, J.C., Introductory Remarks (abstract), EOS, Trans. A.G.U., 57, 907, 1976.
- Cain, J.C., W.E. Daniels, S.J. Hendricks, and D.C. Jensen, An Evaluation of the Main Geomagnetic Field, 1940-1962, J. Geophys. Res., 70, 3647-3674, 1965.
- Cain, J.C., S.J. Hendricks, R.A. Langel, and W.V. Hudson, A Proposed Model for the International Geomagnetic Reference Field - 1965, J. Geomagnet. Geoelec., Kyoto, 19, 335-355, 1967.
- Harwood, J.M., and S.R.C. Malin, Present Trends in the Earth's Magnetic Field, Nature, 259, 469-471, 1976.
- IAGA Division 1 Study Group, International Geomagnetic Reference Field 1975, EOS, Trans. A.G.U., 57, 120-121, 1976.
- Langel, R.A., Near-Earth Magnetic Disturbance in Total Field at High Latitudes 1. Summary of Data from OGO 2, 4, and 6, J. Geophys. Res., 79, 2363-2371, 1974.
- Langel, R.A., Effects of Orbit Error on Satellite Magnetic Field Experiments, GSFC Report X-922-76-124, June 1976.

- Langel, R.A., Near-Earth Satellite Magnetic Field Measurements; A Prelude to Magsat, EOS, Trans. A.G.U., 60, 667-668, 1979.
- Langel, R.A., and J.H. Berbert, Magsat Data Management Plan, GSFC Report, July 1979.
- Langel, R.A., R.D. Regan, and J.P. Murphy, Magsat: A Satellite for Measuring Near-Earth Magnetic Fields, GSFC Report X-922-77-199, July 1977.
- Langel, R.A., R. Coles, and M.A. Mayhew, Comparison of Magnetic Anomalies of Lithospheric Origin Measured by Satellite and Airborne Magnetometers over Western Canada, Can. J. Earth Sci., in press, 1980.
- Leaton, B.R., S.R.C. Malin, and M.J. Evans, An Analytical Representation of the Estimated Geomagnetic Field and Its Secular Change for the Epoch 1965.0, J. Geomagn. Geoelectr., Kyoto, 17, 187-194, 1965.
- Malin, S.R.C., Geomagnetic Secular Variation and its Changes, 1942.5 to 1962.5, Geophys. J.R. Astr. Soc., 17, 415-441, 1969.
- Mead, G.D., An Evaluation of Recent Field Models, in Quantitative Modeling of Magnetospheric Processes, W.P. Olson, Editor, Geophysical Monograph 21, 110-117, American Geophysical Union, 1979.
- NASA/GSFC, Magsat Magnetic Field Satellite, Resource Observation Program, Technical/Descriptive Document, 1979.
- Peddie, N.W., and E.B. Fabiano, A Model of the Geomagnetic Field for 1975, J. Geophys. Res., 81, 2539-2542, 1976.
- Zmuda, A.J., editor, World Magnetic Survey 1957-1969, IAGA Bulletin No. 28, 1971.

Table 1. The MGST(3/80) field model. The mean radius of the Earth was assumed to be 6371.2 km. Mean epoch is 1979.85.

n	m	$s_n^m, \text{nT}$	$h_n^m, \text{nT}$	n	m	$s_n^m, \text{nT}$	$h_n^m, \text{nT}$
1	0	-29990.1	0.0	9	8	1.5	-6.7
1	1	-1958.1	5609.5	9	9	-4.5	3.5
2	0	-1993.2	0.0	10	0	-3.1	0.0
2	1	3027.7	-2129.5	10	1	-3.6	1.3
2	2	1662.4	-192.3	10	2	2.4	0.6
3	0	1269.6	0.0	10	3	-5.2	2.5
3	1	-2180.1	-331.4	10	4	-2.0	5.5
3	2	1251.7	270.6	10	5	4.6	-4.3
3	3	833.3	-250.6	10	6	3.2	-0.1
4	0	937.2	0.0	10	7	0.6	-1.4
4	1	782.7	211.9	10	8	2.1	3.5
4	2	399.2	-257.6	10	9	3.5	-0.4
4	3	-419.8	51.3	10	10	-1.0	-6.5
4	4	198.7	-297.6	11	0	2.2	0.0
5	0	-213.7	0.0	11	1	-1.3	0.7
5	1	357.1	42.9	11	2	-1.8	1.9
5	2	261.0	148.8	11	3	2.2	-1.1
5	3	-73.3	-150.5	11	4	0.2	-2.7
5	4	-162.8	-78.8	11	5	-0.5	0.7
5	5	-46.5	90.6	11	6	-0.3	-0.1
6	0	48.9	0.0	11	7	1.9	-2.3
6	1	65.3	-14.4	11	8	1.9	0.0
6	2	41.1	93.8	11	9	-0.3	-1.6
6	3	-192.2	70.9	11	10	2.1	-0.8
6	4	3.9	-43.2	11	11	2.4	0.8
6	5	14.1	-2.7	12	0	-1.7	0.0
6	6	-107.8	17.2	12	1	0.6	0.9
7	0	70.6	0.0	12	2	-0.1	0.3
7	1	-58.6	-81.2	12	3	-0.1	2.2
7	2	1.9	-27.2	12	4	0.3	-1.5
7	3	20.5	-5.1	12	5	0.6	0.6
7	4	-12.3	16.4	12	6	-0.6	0.3
7	5	-0.2	18.6	12	7	-0.4	-0.3
7	6	10.9	-22.9	12	8	-0.0	0.3
7	7	-2.1	-9.2	12	9	-0.7	-0.3
8	0	18.1	0.0	12	10	0.0	-1.5
8	1	6.7	6.7	12	11	0.1	-0.5
8	2	-0.2	-17.8	12	12	1.4	-0.3
8	3	-10.8	4.2	13	0	0.1	0.0
8	4	-7.1	-22.1	13	1	-0.4	-0.2
8	5	4.4	9.3	13	2	0.3	0.4
8	6	2.4	16.0	13	3	-0.5	1.7
8	7	6.0	-13.1	13	4	-0.1	0.1
8	8	-1.8	-15.8	13	5	1.2	-0.6
9	0	5.7	0.0	13	6	-0.5	-0.1
9	1	10.1	-21.3	13	7	0.1	1.0
9	2	1.0	15.0	13	8	-0.4	0.1
9	3	-12.8	9.0	13	9	-0.0	0.9
9	4	9.3	-4.9	13	10	0.0	0.4
9	5	-3.1	-6.7	13	11	-0.1	-0.4
9	6	-1.3	8.7	13	12	0.7	-1.1
9	7	6.6	9.5	13	13	0.4	0.5

TABLE 2. STRENGTH OF CENTERED DIPOLE AND RATES OF CHANGE  
FOR THE PERIOD 1960 TO 1980

<u>Model</u>	<u>Epoch</u>	<u><math>MR^{-3}, nT</math></u>	<u><math>\dot{MR}^{-3}, nT/yr</math></u>
Cain et al., 1965	1960	31043	-19
GSFC(12/66) (Cain et al., 1967)	1960	31021	-15
Malin, 1969	1960	31152	-17
Leaton et al., 1965	1965	30987	-16
Barracough et al., 1978	1965	30955	-22
POGO(2/72) (Langel et al., 1980)	1968	30879	-26
IGS/75 (Barracough et al., 1975)	1975	30701	-29
AWC/75 (Peddle and Fabiano, 1976)	1975	30652	-26
MGST(3/80)	1979.85	30573	-26*

\*by comparison with POGO(2/72), Epoch 1968

Table 3: Residuals of selected MAGSAT data to some published field models. Units are nT.

	IGRF 1975	AWC/75	IGS/75	POGO(2/72)	MGST(3/80)
Scalar:mean deviation	-90	61	23	9	0
:standard deviation	125	127	120	107	8
$B_r$ :mean deviation	29	46	40	25	21
:standard deviation	204	153	137	211	44
$B_\theta$ :mean deviation	44	-10	8	12	12
:standard deviation	146	115	114	145	107
$B_\phi$ :mean deviation	62	62	61	61	62
:standard deviation	181	157	155	208	129






Figure 1. Distribution of scalar field data used to derive model. Vector data was limited to  $\pm 50^\circ$  latitude.

ORIGINAL PAGE IS  
OF POOR QUALITY






C) HORIZONTAL FIELD, H, GAUSS.




D) VERTICAL FIELD, Z, GAUSS

Figure 2(c) and 2(d). Horizontal field (c) and vertical field (d)  
at 400 km altitude, as computed from the MGST(3/80) model.

ORIGINAL PAGE IS  
OF POOR QUALITY



**A) FIELD MAGNITUDE, B, GAUSS.**



**B) DECLINATION, D, DEGREES.**

Figure 2. Geomagnetic field magnitude (a) and declination (b) at 400 km altitude, as computed from the MGST(3/80) model.




Figure 3. Strength of the Centered Dipole since 1960. Slope of the line is

-27 nT/year.

# BIBLIOGRAPHIC DATA SHEET

1. Report No. TM 80679	2. Government Accession No.	3. Recipient's Catalog No.	
4. Title and Subtitle  INITIAL GEOMAGNETIC FIELD MODEL FROM MAGSAT		5. Report Date April 1980	
		6. Performing Organization Code 922	
7. Author(s) R.A. Langel, R.H. Estes, G.D. Mead, E.B. Fabiano, E.R. Lancaster		8. Performing Organization Report No.	
9. Performing Organization Name and Address Geophysics Branch Goddard Space Flight Center Greenbelt, MD 20771		10. Work Unit No.	
		11. Contract or Grant No.	
12. Sponsoring Agency Name and Address  Same As Above		13. Type of Report and Period Covered	
		14. Sponsoring Agency Code	
15. Supplementary Notes			
16. Abstract <p>Magsat data from the magnetically quiet days of November 5-6, 1979, were used to derive a thirteenth degree and order spherical harmonic geomagnetic field model MGST(3/80). The model utilized both scalar and vector data and fit that data with standard deviations of 8, 52, 55 and 97 nT for the scalar magnitude, <math>B_r</math>, <math>B_\theta</math>, and <math>B_\phi</math>, respectively. When compared with earlier models, the earth's dipole moment continues to decrease at a rate of about 26 nT/yr. Evaluation of earlier models with Magsat data shows that the scalar field at the Magsat epoch is best predicted by the POGO(2/72) model but that the AWC/75 and IGS/75 are better for predicting vector fields.</p>			
17. Key Words (Selected by Author(s)) Magsat Magnetic field satellite geomagnetic field models		18. Distribution Statement	
19. Security Classif. (of this report)	20. Security Classif. (of this page)	21. No. of Pages	22. Price*